

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number
WO 01/52816 A2

- (51) International Patent Classification⁷: **A61K 9/00**
- (21) International Application Number: **PCT/US01/00817**
- (22) International Filing Date: **11 January 2001 (11.01.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
09/488,406 **19 January 2000 (19.01.2000)** **US**
- (71) Applicant (*for all designated States except US*): **CARLSBAD TECHNOLOGY, INC.** [US/US]; 5923 Balfour Court, Carlsbad, CA 92008 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **LEE, Fang-Yu** [—/—]; 1191, Sec. 1, Chung Shan Road, Taiwan Tachia, Taichung (TW). **KUO, Hang-Chiang** [—/—]; 1191, Sec. 1, Chung Shan Road, Taiwan Tachia, Taichung (TW). **CHEN, Shan-Chiung** [—/—]; 1191, Sec. 1, Chung Shan Road, Tachia, Taichung (TW).
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **ORALLY ADMINISTERED PHARMACEUTICAL FORMULATIONS OF BENZIMIDAZOLE DERIVATIVES AND THE METHOD OF PREPARING THE SAME**

(57) Abstract: The present invention provides pharmaceutical formulations which contain (a) an inert core of sugar, sugar and starch, or microcrystalline cellulose, (b) a drug emulsion layer which is made from mixing a free base of benzimidazole derivative (such as omeprazole or lansprazole) with a nonionic surfactant and water, (c) a protective coating which is made of a film-forming compound, and optionally a plasticizer or excipient, and (d) an enteric coating which is made of a pharmaceutically acceptable polymer and a plasticizer. Optionally, a basic amino acid can be added to the drug emulsion layer or the protective coating. The present invention also provides the method for making the pharmaceutical formulations.

WO 01/52816 A2

**ORALLY ADMINISTERED PHARMACEUTICAL FORMULATIONS
OF BENZIMIDAZOLE DERIVATIVES AND THE METHOD OF
PREPARING THE SAME**

5 **FIELD OF THE INVENTION**

This invention relates to novel orally administered pharmaceutical formulations in the form of granules which comprises, as an active ingredient, a potent gastric acid secretion inhibitor, i.e., a substituted 2-(2-benzimidazolyl)-pyridine such as omeprazole or lansoprazole, and the process of making the formulations.

10

BACKGROUND OF THE INVENTION

Benzimidazole derivatives have been known for their anti-ulcer activities as inhibitors of gastric acid secretion. For example, omeprazole, which has the formula of 5-methoxy-2(((4-methoxy-3,5-dimethyl-2-pyridinyl)methyl)sulfinyl)-1H-benzimidazole), is known for its
15 activity as an inhibitor of H^+K^+ -ATPase and the proton pump in the gastric mucosa and can be used for the treatment of gastric and duodenal ulcers (Pilbrant and Cederberg, Scand. J. Gastroenterology (1985) 20:113-120). The information of omeprazole can be found U.S. Pat. No. 4,255,431, U.S. Pat. No. 4,786,505, and EPO 124495. Lansoprazole, which has the formula of 2-[[[3-methyl-4-(2,2,2-trifluoroethoxy)-2-pyridyl]methyl] sulfinyl]benzimidazole,
20 is useful for prophylaxis and therapy of digestive ulcers (e.g., gastric ulcer, duodenal ulcer) and gastritis. Its empirical formula is $C_{16}H_{14}F_3N_3O_2S$ with a molecular weight of 369.37. The information of lansoprazole can be found U.S. Pat. No. 4,628,098, U.S. Pat. No. 4,689,333, and U.S. Pat. No. 5,026,560.

Omeprazole is very slightly soluble in water, but very soluble in alkaline solutions as
25 the negatively charged ion. It is an ampholyte with $pK_a \sim 4$ (pyridinium) and 8.8

5 (benzimidazole). Lansoprazole is a white to brownish-white odorless crystalline powder which melts with decomposition at approximately 166°C. Lansoprazole is freely soluble in dimethylformamide; soluble in methanol; sparingly soluble in ethanol; slightly soluble in ethyl acetate, dichloromethane and acetonitrile; very slightly soluble in ether; and practically insoluble in hexane and water.

10 According to Pilbrant and Cederberg, Scand. J. Gastroenterology (1985) 20:113-120, omeprazole is susceptible to degradation/-transformation in acid and neutral media. The rate of degradation proceeds with a half-life of less than 10 minutes at pH-values below 4. At pH 6.5 the half-life of degradation is 18 hours; at pH 11 about 300 days.

Due to the acidic gastric condition, a pharmaceutical dosage form of omeprazole must
15 be coated with an enteric coating to prevent omeprazole from premature contact with gastric juice. However, ordinary enteric coatings are also made of acidic compounds. Therefore, if omeprazole is directly covered with the conventional enteric coating, the dosage form may not only badly become discolored but also decreased in omeprazole content with the passage of time. To overcome the acidic labile problem and to prolong the storage stability of
20 omeprazole, it is generally recommended to mix omeprazole with an alkaline material so as to create a high pH value for the drug. For instance, U.S. Pat. No. 4,738,974 describes the alkaline salts of omeprazole which include Li^+ , Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Ti^{4+} , $\text{N}^+(\text{R}^1)_4$ or guanidinium salts.

Alternatively, U.S. Pat. No. 4,786,505 describes an oral dosage form of omeprazole,
25 where omeprazole is mixed with an alkaline reacting substance to create a "micro-pH" around each omeprazole particle of not less than pH=7, preferably not less than pH=8. The alkaline substances described in U.S. Pat. No. 4,786,505 include the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric acid, carbonic acid, citric acid or other suitable

5 weak inorganic or organic acids; substances normally used in antacid preparations such as aluminium, calcium and magnesium hydroxides; magnesium oxide or composite substances, such as $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$, $(\text{Mg}_6\text{Al}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O})$, $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ or similar compounds; organic pH-buffering substances such as trihydroxymethylaminomethane or other similar pH-buffering substances. The high pH-value of omeprazole can be achieved
10 by using an alkaline reacting salt of omeprazole as described in U.S. Pat. No. 4,738,974.

U.S. Pat. No. 5,232,706 describes an oral dosage form of omeprazole which contains a nucleus formed by a mixture of omeprazole or an alkali salt of omeprazole with a first basic compound, a first coating which contains at least an excipient and a second basic compound, and an enteric coating. The basic compounds referred to in U.S. Pat. No. 5,232,706 are
15 sodium, potassium, magnesium, calcium, aluminum or dihydroxyaluminium salts of amino acids, such as glycocoll, glutamic acid, or lysine, or a pyridine carboxylic acids such as nicotinic acid, or organic bases such as guanidine.

Due to the insolubility of omeprazole in water, most of the omeprazole formulations are prepared by mixing the powder form of omeprazole with various kinds of binders, excipients
20 and carriers. For example, U.S. Pat. No. 4,786,505 describes the preparation of the omeprazole core by mixing omeprazole with alkaline reacting substances to form a powder mixture, followed by formulating the powder mixture into small beads, i.e., pellets, tablets, hard gelatine or soft gelatine capsules by conventional pharmaceutical procedures, i.e., by pressing through an extruder and spheronized to pellets.

25 U.S. Pat. No. 5,385,739 discloses omeprazole microgranules where the powder form of omeprazole is diluted with a substantially equal amount of mannitol powder, together with sodium lauryl sulfate and carboxymethylstarch, so as to produce a homogeneous and stable dilute powder.

5 U.S. Pat. No. 5,026,560 describes a formulation of making spherical granules containing omeprazole or lansoprazole. The formulation contains a spherical granule which has a core coated with a binder and spraying powder containing the drug and low substituted hydroxypropylcellulose.

10 In the invention to be described, novel orally administered pharmaceutical formulations of omeprazole or lansoprazole will be described. These formulations are distinctively different from those of the patents described above: First, the invention uses a free base of omeprazole or lansoprazole instead of the alkaline salt form of the drug. Second, the free base of omeprazole or lansoprazole is mixed with a non-ionic surfactant and water to form an emulsion, rather than a powder mix, which then can be sprayed and dried onto an inert core to
15 form a granule. Third, the invention demonstrates that it is not necessary to mix the free base of omeprazole or lansoprazole with any alkaline substance in order to create a fully bioavailable dosage form. In fact, the omeprazole formulations in which omeprazole is not mixed with any alkaline substance display equal or better dissolution rate than the commercially available omeprazole formulation such as prilosec where omeprazole is mixed
20 with an alkaline substance.

SUMMARY OF THE INVENTION

A first embodiment of the present invention provides an orally administered pharmaceutical granule of omeprazole or lansoprazole which contains (a) an inert core which is
25 made of starch, a mixture of sugar and starch, or microcrystalline cellulose; (b) a drug emulsion deposited on the inert core, wherein said drug emulsion comprises an effective amount of a free base of omeprazole or lansoprazole, a non-ionic surfactant, a basic amino acid, and water; (c) a protective coating deposited on top of the drug coating, wherein the

5 protective coating comprises at least one film-forming compound which is selected from the group consisting of hydroxypropyl methylcellulose (HPMC), hydroxypropyl cellulose (HPC), polyvinyl pyrrolidone (PVP) and hydroxymethylcellulose (HMC), optionally a plasticizer; and (d) an enteric coating deposited on top of the protective coating, wherein the enteric coating comprises at least a pharmaceutically acceptable polymer which is selected from the group
10 consisting of hydroxypropylmethyl-cellulose phthalate (HPMCP), cellulose acetate phthalate (CAP), co-polymerized methacrylic acid/methacrylic acid methyl esters, and the plasticizer. The pharmaceutical granules can be encapsulated. The granules can also be compressed into tablets by mixing the granule with at least one excipient which is selected from the group consisting of lactose, starch, talc, microcrystalline cellulose, and polyethylene glycol (PEG).

15 Preferably, the drug emulsion contains 30-60 wt% of water, 1-10wt% of non-ionic surfactant, and 1-15wt% of basic amino acid. The basic amino acid is used to stabilize the drug and to prevent the drug from decoloration with passage of time. The preferable basic amino acid used for this purpose includes arginine, lysine, histidine, and tryptophane. The most favorable one is arginine. The preferable nonionic surfactant used in the drug emulsion is
20 polyoxypropylene-polyoxyethylene block copolymers (e.g., Poloxamer 188) or polyethylene-sorbitan-fatty acid esters (e.g., Tween 80), most favorably Poloxamer 188. The preferable plasticizer used in the protective coating and enteric coating include triethyl citrate, triacetin, and diethyl phthalate, most favorably triethyl citrate. The weight ratio of the polymer and plasticizer in the enteric coating is preferably no less than 10:1, and more preferably no less
25 than 50:1.

The pharmaceutical granule is made by the following steps: (a) preparing an inert core; (b) coating the inert core with a drug emulsion which comprises a free base of omeprazole or lansoprazole, a non-ionic surfactant, a basic amino acid, and water by spraying the drug

5 emulsion onto the inert core which then can be dried using a conventional way, e.g., under warm air; (c) coating the drug emulsion (after drying) with a protective coating which comprises a film-forming compound, a plasticizer, and water by spraying the protective coating onto the dried drug emulsion followed by drying; (d) covering the protective coating with an enteric coating which comprises a polymer, a plasticizer, and water by spraying the enteric
10 coating onto the protective coating.

This pharmaceutical formulation can be used to treat patients with gastro-intestinal disorders, including ulcers (e.g., gastric and duodenal ulcers). The treatment includes orally administering a therapeutically effective amount of the formulation to a host in need of such treatment.

15 The second embodiment of the present invention provides an oral administered pharmaceutical granule of omeprazole or lansoprazole which contains: (a) an inert core which is made of starch, a mixture of sugar and starch, or microcrystalline cellulose; (b) a drug emulsion deposited on the inert core, wherein the drug emulsion comprises an effective amount of a free base of omeprazole or lansoprazole, a non-ionic surfactant, and water, wherein the
20 drug emulsion does not contain an alkaline salt or compound; (c) a protective coating deposited on the drug emulsion, wherein the protective coating comprises at least one film-forming compound which is selected from the group consisting of hydroxypropyl methylcellulose (HPMC), hydroxypropyl cellulose (HPC), polyvinyl pyrrolidone (PVP) and hydroxy methylcellulose, and optionally polyethylene glycol (PEG); and (d) an enteric coating
25 deposited on the protective coating, wherein the enteric coating comprises at least a polymer which is selected from the group consisting of hydroxypropylmethylcellulose phthalate (HPMCP), cellulose acetate phthalate (CAP), and co-polymerized methacrylic acid/methacrylic acid methyl esters, and a plasticizer. The pharmaceutical granules can be encapsulated. The

5 granules can also be compressed into tablets by mixing the granule with at least one excipient which is selected from the group consisting of lactose, starch, talc, microcrystalline cellulose, and polyethylene glycol (PEG).

Preferably, the nonionic surfactant in the drug emulsion is polyoxypropylene-polyoxyethylene block copolymers (e.g., Poloxamer 188) or polyethylene-sorbitan-fatty acid
10 esters (e.g., Tween 80), most favorably Poloxamer 188. Also preferably, the protective coating contains a basic amino acid, e.g., arginine, lysine, histidine, or tryptophane. The most preferable basic amino acid is arginine. The addition of a basic amino acid in the protective coating not only provide better stability for the formulation but also help to prevent the drug from decoloration with the passage of time. PEG is a preferable excipient used in the
15 protective coating. The most preferable PEG is PEG 6000 which has a molecular weight range between 7000 and 9000. The plasticizer used in the enteric coating includes triethyl citrate, triacetin, and diethyl phthalate, preferably triethyl citrate. The weight ratio of the polymer and plasticizer in the enteric coating is preferably no less than 10:1, and more preferably no less than 50:1. In addition, the protective coating can have one or more sublayers. It is preferably
20 that at least one of the sublayers contains a basic amino acid.

The process for making the pharmaceutical granule described in the second embodiment includes the following steps: (a) preparing an inert core; (b) coating the inert core with a drug emulsion which comprises a free base of omeprazole or lansoprazole, a non-ionic surfactant, and water by spraying the drug emulsion onto the inert core, wherein the drug
25 emulsion is free of any alkaline salt or compound; (c) coating the drug emulsion with a protective coating which comprises a film-forming compound, an excipient (e.g., PEG 6000), and water by spraying the protective coating onto the drug emulsion; and (d) covering the

5 protective coating with an enteric coating which comprises a polymer, a plasticizer, and water by spraying said enteric coating onto said protective coating.

The formulation shown in the second embodiment can be used to treat gastro-intestinal disorders, including ulcers (e.g., gastric and duodenal ulcers). The method for treating gastro-intestinal disorders comprise orally administered a therapeutically effective amount of the drug
10 method for treating ulcer comprising orally administering to a host in need of such treatment a therapeutically effective amount of the formulation.

DETAILED DESCRIPTION OF THE INVENTION

The pharmaceutical granule described in the present invention contains four distinctive
15 layers, which are: (1) an inert core, (2) a drug emulsion, (3) a protective coating, and (4) an enteric coating.

The inert core is made of sucrose, starch, talc, or microcrystalline cellulose, alone or in any combination, and is free of any alkaline salt or compound. The inert core can be produced by a granulator or bought from any bulk drug manufacturers. There are three major kinds of
20 inert cores which are commercially available, which are: (1) pure sugar cores; (2) cores containing a mixture of sugar and starch; and (3) microcrystalline cores.

The drug emulsion is produced by mixing the drug, e.g., omeprazole or lansoprazole, with a non-ionic surfactant, e.g., Poloxamer 188 or Tween 80, and water, to form a homogeneous emulsion. Poloxamer 188 is a polyoxypropylene-polyoxyethylene block
25 copolymer having an average molecular weight of 8350. Tween 80 is the trademark for polyethylene-sorbitan-fatty acid esters. Optionally, a basic amino acid, such as arginine, lysine, histidine, and tryptophane, can be added to the drug emulsion. The addition of the basic amino acid in the drug emulsion helps to prevent the drug from decoloration. The drug emulsion has

5 the following advantages over the prior art references disclosed in the "Background of the Invention" section: (1) no binder is needed (in the prior art references, a binder is required because the drug is in the powdery form); (2) no alkaline salt or compound is needed to be mixed with the drug and the drug emulsion helps to preserve the stability and content of the drug with passage of time; and (3) the drug is homogeneously suspended in the emulsion
10 which then can be quantitatively and evenly distributed onto the inert core. The drug emulsion can be coated onto the inert core by conventional coating procedures or sprayed onto the inert core by use of a fluidized bed coating device such as the Glatt or Huttlin machine.

The protective coating serves to separate the drug from the enteric coating. It can contain more than one sublayers. The material for the separating layer is chosen among the
15 pharmaceutically acceptable, water soluble polymers used for film-coating applications, such as hydroxypropylmethyl cellulose (HPMC), hydroxypropyl cellulose (HPC), polyvinyl pyrrolidone (PVP), or hydroxymethyl cellulose (HMC). Optionally, a plasticizer can be added to the protective coating. The plasticizer which can be used in the protective coating includes diethyl phthalate, triacetin, and triethyl citrate. Also optionally, isopropyl alcohol and
20 methylene chloride could be used to replace purified water in the protective coating. In addition, in the protective coating of the second embodiment, optionally an excipient such as PEG 6000 is used to replace the plasticizer. The protective coating can be coated onto the inert core by conventional coating procedures or sprayed onto the inert core by use of a fluidized bed coating device such as the Glatt or Huttlin machine.

25 The enteric coating serves to protect the drug from contact with acidic gastric juice. The material used in the enteric coating includes Eudragit L, Eudragit S (Rohm Pharma), hydroxypropyl methylcellulose phthalate (HPMCP), or cellulose acetate phthalate (CAP). Eudragit L and Eudragit S are tradenames of co-polymerized methacrylic acid/methacrylic acid

5 methyl esters. The enteric coating also contain a plasticizer such as diethyl phthalate, triethyl citrate, or triacetin. The weight ratio of the polymer to plasticizer should be no less than 10:1, preferably no less than 50:1. The enteric coating is applied onto the protective layer by conventional coating procedures or sprayed onto the inert core by use of a fluidized bed coating device such as the Glatt or Huttlin machine.

10 The invention is described in details in the following examples:

EXAMPLE 1:

A. Formulation of the Pharmaceutical Granules:

(1) Inert Core: 1097.6 g of sugar, sugar plus starch (in any combinations), or microcrystalline cellulose.

15 (2) Drug Emulsion:

Omeprazole	147 g
Poloxamer 188	98 g
Arginine	78.4 g
Purified Water	924 ml

20 (3) Protective Coating:

HPMC	78.4 g
Triethyl Citrate	7.84 g
Purified Water	784 ml

(4) Enteric Coating Layer:

25 Eudragit L30D	1437.33 g
Triethyl Citrate	21.56 g
Purified Water	478.8 ml

B. Method of Preparing the Formulation:

The inert core was either bought from companies selling bulk drug material or prepared using the Glatt machine (a fluidized bed particle making machine imported from Germany).

30

5 The drug emulsion was prepared by mixing omeprazole, Poloxamer 188, and
arginine in purified water. The emulsion was then placed into the spray gun of the Glatt
machine and sprayed onto the core particles while the Glatt machine was set in running
(circulating) condition. This would allow the drug to be evenly coated onto the core
particles to form drug-coated spherical particles. The drug-coated particles were dried
10 under warm air within the Glatt machine.

 Then, a protective coating solution was prepared by mixing HPMC and triethyl
citrate in purified water. This coating was then placed into the spray gun of the Glatt
machine and sprayed onto the drug-coated particles while the Glatt machine was set in
running condition. After the coating was completed, the protective coating-covered
15 particles were again dried under warm air within the Glatt machine.

 Finally, an enteric coating was prepared by mixing Eudragit L30D and triethyl
citrate in purified water. This coating was placed into the spray gun of the Glatt
machine and sprayed onto the protective coating-covered particles to form the
pharmaceutical granules before final drying of the granules to complete the process of
20 making the enteric coating-covered granules.

EXAMPLE 2:

A. Formulation of Pharmaceutical Granules:

- (1) Inert Core: 1187.41 g of sugar, sugar plus starch (in any combinations), or
microcrystalline cellulose.
- 25 (2) Drug Emulsion:
- | | |
|----------------|-------------------|
| Lansoprazole | 149.8 (or more) g |
| Poloxamer 188 | 98 g |
| Arginine | 78.4 g |
| Purified Water | 910 ml |

5	(3)	<u>Protective Coating:</u>	
		HPMC	78.4 g
		Triethyl Citrate	7.84 g
		Purified Water	784 ml
	(4)	<u>Enteric Coating Layer:</u>	
10		Eudragit L30D	1143.3 g
		Triethyl Citrate	17.15 g
		Purified Water	378 ml

B. Method of Preparing the Formulation:

Same as EXAMPLE 1.

15 EXAMPLE 3:

A. Formulation of Pharmaceutical Granules:

	(1)	<u>Inert Core:</u>	1263.15 g of sugar, sugar plus starch (in any combinations), or microcrystalline cellulose.
	(2)	<u>Drug Emulsion:</u>	
20		Omeprazole	157.5 g
		Poloxamer 188	126 g
		Arginine	84 g
		Purified Water	855 ml
	(3)	<u>Protective Coating:</u>	
25		HPMC	126 g
		Triethyl Citrate	12.6 g
		Isopropyl Alcohol	1764 ml
		Methylene Chloride	756 ml
	(4)	<u>Enteric Coating Layer:</u>	
30		Eudragit L30D	1050 g
		Triethyl Citrate	15.75 g
		Purified Water	345 ml

5 B. Method of Preparing the Formulation:

Same as EXAMPLE 1.

Tests of the Stability of the Pharmaceutical Formulations as Shown in EXAMPLES 1-3

Table I shows the results of the stability studies of the pharmaceutical formulations as
 10 described in EXAMPLES 1-3. The pharmaceutical granules were shelved for 0-6 months at
 75% relative humidity and at 30°C, 37°C and 45°C. A shelf-life of 6 months at 45°C is
 equivalent to a shelf-life of 3 years under normal temperature (25°C).

15 TABLE I.: STABILITY TEST FOR THE PHARMACEUTICAL GRANULES OF
 EXAMPLES 1-3

	Temperature (°C)	Storage Time (Months)	EXAMPLE 1	EXAMPLE 2 (wt%)	EXAMPLE 3
20	Control	0	103.5%	104.1%	104.5%
	30°C	1	103.3%	103.1%	103.9%
	30°C	2	102.7%	102.7%	103.1%
	30°C	3	101.4%	101.9%	102.5%
	30°C	6	97.3%	98.0%	99.7%
25	37°C	1	102.1%	102.7%	102.3%
	37°C	2	101.1%	101.3%	101.5%
	37°C	3	100.1%	100.9%	100.1%
	37°C	6	96.5%	97.2%	98.7%
	45°C	1	101.1%	101.0%	101.6%
30	45°C	2	99.7%	99.3%	99.2%
	45°C	3	97.3%	98.2%	98.7%
	45°C	6	94.8%	96.4%	97.4%

5 The results of Table I indicate that the pharmaceutical granules of Examples 1-3 possess excellent bioavailability and stability at higher temperatures.

 Table II shows the % dissolution, i.e., the percentage of the drug in Examples 1-3 being released in the solution at pH 1.2 (for 120 minutes) and 6.8 (for 30 minutes). The data from the commercially available omeprazole drug (Prilosec[®]) is included for comparison
10 purpose.

TABLE II: % DISSOLUTION OF THE PHARMACEUTICAL GRANULES OF
EXAMPLES 1-3

	pH 1.2	pH 6.8
	(120 minutes)	(30 minutes)
	(% Dissolution)	
EXAMPLE 1	100.6%	88.3%
EXAMPLE 2	100.1%	87.4%
20 EXAMPLE 3	101.2%	90.25%
Prilosec [®]	98.4%	81.4%

 The results of Table II demonstrate that the pharmaceutical granules of Examples 1-3
25 display better release rates (% dissolution) than those of the prior art (Prilosec[®]) formulation.

EXAMPLE 4:

A. Formulation of Pharmaceutical Granules:

- (1) Inert Core: 1294.72 g of sugar, sugar plus starch (in any combinations), or microcrystalline cellulose.
- 30 (2) Drug Emulsion:
- | | |
|----------------|--------|
| Omeprazole | 168 g |
| Poloxamer 188 | 112 g |
| Purified Water | 720 ml |

5	(3)	<u>Protective Coating:</u>	
		HPMC	134.4 g
		PEG 6000	13.44 g
		Purified Water	1680 ml
	(4)	<u>Enteric Coating Layer:</u>	
10		Eudragit L30D	1642.7 g
		Triethyl Citrate	24.64 g
		Purified Water	400 ml

B. Method of Preparing the Formulation:

15 The inert core was either bought from companies selling bulk drug material or prepared using the Glatt machine (a fluidized bed particle making machine imported from Germany).

 The drug emulsion was prepared by first mixing Poloxamer 188 with purified water to form an emulsified solution, followed by adding omeprazole to the solution while stirring to form the drug emulsion. The emulsion was then placed into the spray gun of the Glatt machine and sprayed onto the core particles while the Glatt machine was set in running (circulating) condition. This would allow the drug to be evenly coated onto the core particles to form drug-coated spherical particles. The drug-coated particles were dried under warm air within the Glatt machine.

25 Then, a protective coating solution was prepared by first mixing HPMC with purified water to form a solution, followed by adding PEG 6000 to the solution. This coating was then placed into the spray gun of the Glatt machine and sprayed onto the drug-coated particles while the Glatt machine was set in running condition. After the coating was completed, the protective coating-covered particles were again dried under warm air within the Glatt machine.

30

5 The enteric coating was prepared according to the same procedures as described
in Example 1. This coating was placed into the spray gun of the Glatt machine and
sprayed onto the protective coating-covered particles to form the pharmaceutical
granules before final drying of the granules to complete the process of making the
enteric coating-covered granules.

10 EXAMPLE 5:

A. Formulation of Pharmaceutical Granules:

(1) Inert Core: 1229.76 g of sugar, sugar plus starch (in any combinations), or
microcrystalline cellulose.

(2) Drug Emulsion:

15	Omeprazole	168 g
	Poloxamer 188	89.6 g
	Purified Water	720 ml

(3) Protective Coating:

	HPMC	112 g
20	PEG 6000	11.2 g
	Arginine	112 g
	Purified Water	1723 ml

(4) Enteric Coating Layer:

	Eudragit L30D	1642.7 g
25	Triethyl Citrate	24.64 g
	Purified Water	400 ml

B. Method of Preparing the Formulation:

 The method for preparing the formulation as shown in Example 5 was the same
as that of Example 4 except that in the protective coating, arginine was added to the
30 solution which contained HMPC, PEG 6000 and purified water.

5 EXAMPLE 6:A. Formulation of Pharmaceutical Granules:

(1) Inert Core: 1229.76 g of sugar, sugar plus starch (in any combinations), or microcrystalline cellulose.

(2) Drug Emulsion:

10 Omeprazole 168 g
Poloxamer 188 89.6 g
Purified Water 720 ml

(3) Protective Coating (containing two sublayers):

(i) Sublayer 1 (which is next to the drug emulsion layer)

15 HPMC 44.8 g
PEG 6000 4.48 g
Purified Water 680 ml

(ii) Sublayer 2 (which is next to the enteric coating)

20 HPMC 67.2 g
PEG 6000 6.72 g
Arginine 112 g
Purified Water 1244 ml

(4) Enteric Coating Layer:

25 Eudragit L30D 1642.7 g
Triethyl Citrate 24.64 g
Purified Water 400 ml

B. Method of Preparing the Formulation:

30 The method for preparing the formulation as shown in Example 6 was the same as that of Example 5 except that there were two sublayers within the protective coating which must be applied sequentially, with sublayer 1 coated directly onto the drug coated particles followed by sublayer 2, which then was coated by the enteric coating to form the pharmaceutical granules.

5

Tests of the Stability of the Pharmaceutical Formulations as Shown in EXAMPLES 4-6

Table III shows the % dissolution, i.e., the percentage of the drug in Examples 4-6 being released in the solution at pH 1.2 (for 120 minutes) and at pH 6.8 (for 30 minutes). The data from the commercially available omeprazole drug (Prilosec[®]) is included for comparison purpose.

10

TABLE III.: % DISSOLUTION OF THE PHARMACEUTICAL GRANULES OF
EXAMPLES 4-6

	pH 1.2 (120 minutes)	pH 6.8 (30 minutes)
	(% Dissolution)	
EXAMPLE 4	92.9%	87.3%
EXAMPLE 5	95.4%	86.0%
EXAMPLE 6	100.1%	85.8%
Prilosec [®]	98.4%	81.4%

15

20

25

The results shown in Table III demonstrate that the pharmaceutical formulations of Examples 4-6 display equal or better release rates (% dissolution) than the prior art (Prilosec[®]) formulation.

The stability of the pharmaceutical formulations in Examples 4-6 was also tested for decoloration. The test was conducted under the following conditions: temperature was controlled at 45°C, relative humidity was at 75±5%, storage time was for 1 month. The colors of the pharmaceutical granules in Examples 4-6 were white at the time the granules were prepared.

30

5

TABLE IV.: STABILITY OF THE PHARMACEUTICAL GRANULES AFTER ONE
MONTH OF STORAGE AT 45°C AND 75±5% RELATIVE HUMIDITY

COLOR OF THE GRANULES	
Example 4	Pale Beige
Example 5	White
Example 6	White
Prilosec	Pale Purple

The results of Table IV indicated that while the % dissolution of the pharmaceutical granules in Example 4 was as good as those in Examples 5-6 and the commercially available omeprazole granules Prilosec[®], its stability was probably slightly worse than those of Examples 5-6, although the degree of stability between the granules of Example 4 and Prilosec[®] was probably comparable. Because the major difference between Example 4 and Examples 5-6 was that the protective coating of Example 4 did not contain arginine, it could therefore conclude that the addition of arginine (or other basic amino acid) in the protective coating had contributed to the stability of the pharmaceutical granules.

While the invention has been described by way of examples and in terms of the preferred embodiments, it is to be understood that the invention is not limited to the disclosed embodiments. On the contrary, it is intended to cover various modifications as would be apparent to those skilled in the art. Therefore, the scope of the appended claims should be accorded the broadest interpretation so as to encompass all such modifications.

5 What is claimed is:

1. An orally administered pharmaceutical granule comprising:
an inert core which is made of starch, a mixture of sugar and starch, or
microcrystalline cellulose;
a drug emulsion deposited on said inert core to form a drug coating, wherein said
10 drug emulsion comprises an effective amount of free base omeprazole or lansoprazole, a
non-ionic surfactant, a basic amino acid, and water;
a protective coating deposited on said drug coating, wherein said protective coating
comprises at least one film-forming compound which is selected from the group
consisting of hydroxypropyl methylcellulose (HPMC), hydroxypropyl cellulose (HPC),
15 polyvinyl pyrrolidone (PVP) and hydroxy methylcellulose (HMC), and a plasticizer; and
an enteric coating deposited on said protective coating, wherein said enteric coating
comprises at least a polymer which is selected from the group consisting of
hydroxypropylmethylcellulose phthalate (HPMCP), cellulose acetate phthalate (CAP),
co-polymerized methacrylic acid/methacrylic acid methyl esters, and a plasticizer.
20
2. The orally administered pharmaceutical granule according to claim 1, wherein said
drug emulsion contains 30-60 wt% of water, 1-10 wt% of the non-ionic surfactant, and 1-
15wt % of the basic amino acid.
- 25 3. The orally administered pharmaceutical granule according to claim 1, wherein said
basic amino acid is selected from the group consisting of arginine, lysine, histidine, and
tryptophane.
4. The orally administered pharmaceutical granule according to claim 1, wherein said
30 non-ionic surfactant is a polyoxypropylene-polyoxyethylene block copolymer or a
polyethylene-sorbitan-fatty acid ester.
5. The orally administered pharmaceutical granule according to claim 1, wherein said
35 plasticizer is triethyl citrate, triacetin, or diethyl phthalate.

- 5 6. The orally administered pharmaceutical preparation according to claim 1, wherein said polymer and said plasticizer in said enteric layer are at a weight ratio of no less than 10:1.
7. The orally administered pharmaceutical granule according to claim 1, wherein said
10 granule is further encapsulated.
8. The orally administered pharmaceutical granule according to claim 1, wherein said granule is compressed into tablet by mixing with at least an excipient which is selected from the group consisting of lactose, starch, talc, microcrystalline cellulose, and
15 polyethylene glycol (PEG).
9. A process of making an orally administered pharmaceutical granule according to claim 1 comprising:
preparing an inert core;
20 coating the inert core with a drug emulsion which comprises a free base of omeprazole or lansoprazole, a non-ionic surfactant, arginine, and water by spraying said drug emulsion onto said inert core;
coating the drug emulsion with a protective coating which comprises a film-forming compound, a plasticizer, and water by spraying said protective coating onto
25 said drug emulsion;
coating said protective coating with an enteric coating which comprises a polymer, a plasticizer, and water by spraying said enteric coating onto said protective coating.
- 30 10. A method for treating gastro-intestinal disorders comprising orally administering to a host in need of such treatment a therapeutically effective amount of the pharmaceutical granule according to claim 1.

- 5 11. An orally administered pharmaceutical granule comprising:
- an inert core which is made of starch, a mixture of sugar and starch, or microcrystalline cellulose, wherein said inert core is free of alkaline salt or compound;
- a drug emulsion deposited on said inert core to form a drug coating, wherein said drug emulsion comprises an effective amount of free base omeprazole or lansoprazole, a
- 10 non-ionic surfactant, and water, wherein said drug emulsion does not contain an alkaline salt or compound;
- a protective coating deposited on said drug coating, wherein said protective coating comprises at least one film-forming compound which is selected from the group consisting of hydroxypropyl methylcellulose (HPMC), hydroxypropyl cellulose (HPC),
- 15 polyvinyl pyrrolidone (PVP) and hydroxy methylcellulose, optionally an excipient;
- an enteric coating deposited on said protective coating, wherein said enteric coating comprises at least a polymer which is selected from the group consisting of hydroxypropylmethylcellulose phthalate (HPMCP), cellulose acetate phthalate (CAP), and co-polymerized methacrylic acid/methacrylic acid methyl esters, and a plasticizer.
- 20
12. The orally administered pharmaceutical preparation according to claim 11, wherein said non-ionic surfactant is a polyoxypropylene-polyoxyethylene block copolymer or a polyethylene-sorbitan-fatty acid ester.
- 25 13. The orally administered pharmaceutical granule according to claim 11, wherein said excipient is polyethylene glycol 6000.

- 5 14. The orally administered pharmaceutical granule according to claim 11, wherein said polymer and said plasticizer in said enteric layer are at a weight ratio of no less than 10:1.
- 10 15. The orally administered pharmaceutical granule according to claim 11, wherein said protective coating further comprises a basic amino acid selected from the group consisting of arginine, lysine, histidine, and tryptophan.
16. The orally administered pharmaceutical granule according to claim 11, wherein said protective coating comprises one or more sublayers.
- 15 17. The orally administered pharmaceutical granule according to claim 16, wherein at least one of the sublayers contains a basic amino acid selected from the group consisting of arginine, lysine, histidine, and tryptophan.
- 20 18. The orally administered pharmaceutical granule according to claim 16, wherein none of the sublayers of the protective coating contains an alkaline salt or compound.
19. The orally administered pharmaceutical granule according to claim 11, wherein said granule is encapsulated.

- 5 20. The orally administered pharmaceutical granule according to claim 11, wherein
said granule is compressed into tablet by mixing with at least an excipient which is
selected from the group consisting of lactose, starch, talc, microcrystalline cellulose, and
polyethylene glycol (PEG).
- 10 21. A process of making an orally administered pharmaceutical granule according to
claim 11 comprising:
preparing an inert core, wherein said inert core is free of alkaline salt or compound;
coating the inert core with a drug emulsion which comprises a free base of
omeprazole or lansoprazole, a non-ionic surfactant, and water by spraying said drug
15 emulsion onto said inert core, wherein said drug emulsion does not contain an alkaline
salt or compound;
coating the drug emulsion with a protective coating which comprises a film-
forming compound, a plasticizer, and water by spraying said protective coating onto said
drug emulsion;
20 coating the protective layer with an enteric coating which comprises a polymer, a
plasticizer, and water by spraying said enteric coating onto said protective coating.
22. A method for treating gastro-intestinal disorders comprising orally administering to
a host in need of such treatment a therapeutically effective amount of the pharmaceutical
25 granule according to claim 11.